

Copy of search for 09/109063

\*\*\*\*\*

\* Welcome to MESSENGER (APS Text) at USPTO \*

\* The USPTO production files are current through: \*

\* DECEMBER 15, 1998 for U.S. Patent Text Data. \*

\* DECEMBER 15, 1998 for U.S. Current Classification Data. \*

\* DECEMBER 15, 1998 for U.S. Patent Image Data. \*

FILE 'JPO' ENTERED AT 10:22:45 ON 16 DEC 1998

\*\*\*\*\*

\* G P I \*

\* J A P A N E S E P A T E N T A B S T R A C T S \*

\* THE FILE IS CURRENT THROUGH AUGUST 31, 1998. \*

\*\*\*\*\*

L1 210 TRANSGLUTAMINASE

L2 1764 HIROSHI?/IN

L3 0 L1 AND L2

L4 22525 TAKAGI?/IN

L5 4 L1 AND L4

JP4063077A

L5: 1 of 4

TITLE: PRODUCTION OF BACTERIAL \*\*TRANSGLUTAMINASE\*\* AND RELATED MATERIAL

JP405199883A

L5: 2 of 4

TITLE: RECOMBINANT \*\*TRANSGLUTAMINASE\*\*

JP402100647A

L5: 3 of 4

TITLE: PRODUCTION OF FRIED BEAN CURD

JP402069146A

L5: 4 of 4

TITLE: PRODUCTION OF PROTEIN-TEXTURED PRODUCT

JP40630771A

Feb. 8, 1994

L5: 1 of 4

PRODUCTION OF BACTERIAL \*\*TRANSGLUTAMINASE\*\* AND RELATED MATERIAL

INVENTOR: KAWAI, MISAKO TAKEHANA, SHINO \*\*TAKAGI, HIROSHI\*\*

APPLICANT: AJINOMOTO CO INC

APPL NO: JP 04187038

DATE FILED: Jul. 14, 1992 INT-CL: C12N9/10; C12N15/62

ABSTRACT:

PURPOSE: To mass produce bacterial \*\*transglutaminase\*\* by using a bacterium such as *Escherichia coli*.

CONSTITUTION: *Escherichia coli* retaining a plasmid manifesting a fused protein containing bacterial \*\*transglutaminase\*\* and a hydrophilic peptide existing at the amino end side of the bacterial \*\*transglutaminase\*\* is cultured to produce a bacterial

\*\*transglutaminase\*\* fused protein as an inert sealed material in cells. The sealed material is recovered from the cells, the sealed material is solubilized by using a denaturing agent and the denaturing agent is removed to give a fused protein having

\*\*transglutaminase\*\* activity. COPYRIGHT: (C)1994, JPO&lapo

\*\*\*\*\*

JP405199883A

Aug. 10, 1993

L5: 2 of 4

RECOMBINANT \*\*TRANSGLUTAMINASE\*\*

INVENTOR: \*\*TAKAGI, HIROSHI\*\* ARAFUKA, SHINO MATSUI, YUTAKA WASHIZU, KINYA ANDO, KEIICHI

KOIKEDA, SATOSHI

APPLICANT: AMANO PHARMACEUT CO LTD AJINOMOTO CO INC

APPL NO: JP 03267860 DATE FILED: Oct. 16, 1991

INT-CL: C12N15/54; C12N1/19; C12N1/21; C12N9/10; C12N15/70 ADDITIONAL-INT-CL: C07K13/00

ABSTRACT:

PURPOSE: To provide a new DNA useful for producing \*\*transglutaminase\*\*.

CONSTITUTION: A DNA containing a base sequence coding the amino acid sequence of the formula. The DNA is obtained by using as a probe a DNA fragment obtained by PCR method and cloning from the genome DNA of an *Actinomyces* bacterium,

etc. COPYRIGHT: (C)1993, JPO&lapo

(FILE 'HOME' ENTERED AT 12:54:02 ON 21 DEC 1998)

FILE 'REGISTRY' ENTERED AT 12:54:44 ON 21 DEC 1998

L1 4 S SDDRVT/SGSP

FILE 'CAPLUS' ENTERED AT 12:55:16 ON 21 DEC 1998

L2 5 S L1

FILE 'REGISTRY' ENTERED AT 12:57:15 ON 21 DEC 1998

L3 66 S TRANSGLUTAMINASE

FILE 'CAPLUS' ENTERED AT 12:57:29 ON 21 DEC 1998

L4 1938 S L3

L5 356695 S CLON? OR GENE!

L6 208 S L4 AND L5

L7 217 S CODON (5N) (SUBSTITUT? OR OPTIMIZ?)

L8 1 S L4 AND L7

L2 ANSWER 1 OF 5 CAPLUS COPYRIGHT 1998 ACS AN 1994:528753 CAPLUS DN 121:128753

T1 Preparation of bacterial transglutaminase with *Escherichia coli*

IN Kawai, Misako; Takehana, Shino; Takagi, Hiroshi

PA Ajinomoto Kk, Japan

SO Jpn. Kokai Tokkyo Koho, 13 pp. CODEN: JKKXAF DT Patent

LA Japanese FAN CNT 1 PATENT NO. KIND DATE APPLICATION NO. DATE

JP 06030771 A2 19940208 JP 92-187038 19920714

AB A method for the prodn. of bacterial transglutaminase in a microbial host such as *Escherichia coli* is described.

Transglutaminase of *Streptovorticillum* was modified by substitution with hydrophilic amino acid residues to improve its soly. and expressed in *Escherichia coli* as a fusion protein with, e.g., T7 gene 10 peptide. The fusion protein produced in the inclusion bodies was solubilized with a denaturant and cleaved with Factor Xa. The yield of transglutaminase by the method was approx.

20 mg/L.

L2 ANSWER 2 OF 5 CAPLUS COPYRIGHT 1998 ACS AN 1994:209639 CAPLUS DN 120:209639

T1 Chemical synthesis of the gene for microbial transglutaminase from *Streptovorticillum* and its expression in *Escherichia coli*

AU Takehana, Shino; Washizu, Kinya; Ando, Keiichi; Koikeda, Satoshi; Takeuchi, Kazuyuki; Matsui, Hiroshi; Motoki, Masao; Takagi, Hiroshi

CS Food Res. Dev. Lab., Ajinomoto Co., Inc., Kawasaki, 210, Japan

SO Biosci. Biotechnol., Biochem. (1994), 58(1), 88-92 CODEN: BBBIEJ, ISSN: 0916-8451

DT Journal LA English

AB The gene coding for microbial transglutaminase (T Gase) from *Streptovorticillum*, which consists of 331 amino acids, was chem. synthesized. The codons have been substituted for those mainly favored in yeast. The authors strategy involved the

construction of the T Gase gene in five sections (54 oligomers) that contained unique restriction enzyme sites at both ends, which could readily be ligated to form the full-length product. The chem. synthesized gene was inserted downstream from the ompA

signal peptide of the E. coli expression vector, pJM1-ill-ompA, which carries lpp and lac promoters. The resultant plasmid directed the expression of T Gase, with the activity being secreted mainly into the periplasmic space of E. coli. The induced gene product

was identical with native T Gase in size and in immunol. properties, though the enzyme activity was low.

L2 ANSWER 3 OF 5 CAPLUS COPYRIGHT 1998 ACS AN 1994:209638 CAPLUS DN 120:209638

T1 Molecular cloning of the gene for microbial transglutaminase from *Streptovorticillum* and its expression in *Streptomyces lividans*

AU Washizu, Kinya; Ando, Keiichi; Koikeda, Satoshi; Hirose, Susumu; Matsura, Akira; Takagi, Hiroshi; Motoki, Masao; Takeuchi, Kazuyuki

CS Tsukuba Res. Lab., Amano Pharm. Co., Ltd., Tsukuba, 305, Japan

SO Biosci. Biotechnol., Biochem. (1994), 58(1), 82-7 CODEN: BBBIEJ, ISSN: 0916-8451

DT Journal LA English

AB The microbial transglutaminase (T Gase)-producing strain S-8112 was identified as a variant of *Streptovorticillum mobaraense*. The authors amplified a partial gene fragment by polymerase chain reaction (PCR) using oligonucleotides

synthesized from the amino acid sequence of T Gase, and cloned the gene for T Gase using the PCR amplified fragment as a probe. The gene encoded a precursor of T Gase consisting of 406 amino acid residues, which comprised the prepro region of 75

amino acid residues and the mature region of 331 amino acid residues. The authors expressed the T Gase gene in *Streptomyces lividans* under a tyrosinase promoter, and found an active and mature recombinant enzyme, indicating the

processing of the gene product.

L2 ANSWER 4 OF 5 CAPLUS COPYRIGHT 1998 ACS AN 1993:554777 CAPLUS DN 119:154777

T1 Primary structure of microbial transglutaminase from *Streptovorticillum* sp. strain s-8112

AU Kanaji, Toshiya; Ozaki, Hiroshi; Takao, Toshihumi; Kawajiri, Hideo; Ibe, Hiroyuki; Motoki, Masao; Shimomishi, Yasutsugu  
CS Inst. Protein Res., Osaka Univ., Suita, 565, Japan  
SD J. Biol. Chem. (1993), 268(16), 11565-72 CODEN: JBCHA3; ISSN: 0021-9258  
DT Journal LA English

AB The complete amino acid sequence of transglutaminase (EC 2.3.2.13) (TGase), which is produced by a microorganism, *Streptovorticillum* sp. strain s-8112, and catalyzes the acyl transfer reaction between gamma-carboxyamide groups of glutamine residues in proteins and various primary amines, has been established by a combination of fast atom bombardment mass spectrometry and std. Edman degradn. of peptide fragments produced by treatment of the TGase with various proteolytic enzymes and purified by a reversed-phase high performance liq. chromatog. The TGase consists of 331 amino acid residues with a chem. mol. wt. of 37,863, in agreement with the obsd. mol. wt. (37,869.2 +/- 8.8) dedd. from its electrospray ionization mass spectrum. The sequence of the enzyme is very different from those of mammalian TGases represented by guinea pig liver enzyme. The enzyme contains a sole Cys residue, which is essential for its catalytic activity. Hydrophobic anal. indicated that the secondary structure of the region around the active site Cys residue is similar to those of mammalian TGases. These results suggest that this microbial protein evolved by a different pathway from that of mammalian TGases and acquired acyl transfer activity during the evolutionary process.

L2 ANSWER 5 OF 5 CAPLUS COPYRIGHT 1998 ACS AN 1992:629103 CAPLUS DN 117229103

L6 ANSWER 1 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI GTP-dependent conformational changes associated with the functional switch between G.alpha. and crosslinking activities in brain-derived tissue transglutaminase

L6 ANSWER 2 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Preparation and use of respiratory-deficient cells as expression hosts for the manufacture of foreign proteins

L6 ANSWER 3 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI The human prostate-specific transglutaminase gene (TGAM): genomic organization, tissue-specific expression, and promoter characterization

L6 ANSWER 4 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Genotype/phenotype correlation in autosomal recessive lamellar ichthyosis

L6 ANSWER 5 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Plasma transglutaminase in hypertrophic chondrocytes: expression and cell-specific intracellular activation produce cell death and externalization

L6 ANSWER 6 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Induction of differentiation in normal human keratinocytes by adenovirus-mediated introduction of the  $\alpha 2a$  and  $\alpha 2b$  isoforms of protein kinase C

L6 ANSWER 7 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI A novel insertion mutation (T286insC) in exon 9 of the factor XIII-A subunit gene

L6 ANSWER 8 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Isolation, properties and applications of a thermostable endo-beta-1,4-glucanase

L6 ANSWER 9 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Coexpression of p53 and tissue transglutaminase \*\*\*genes\*\*\* in human normal and pathologic adrenal tissues

L6 ANSWER 10 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI \*\*\*Cloning\*\*\* and expression of heat-stable bacterial endoglycanase gene and use of enzyme in industrial processes

L6 ANSWER 11 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Molecular \*\*\*cloning\*\*\* of the transglutaminase gene from *Bacillus subtilis* and its expression in *Escherichia coli*

L6 ANSWER 12 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Purification, characterization, and gene \*\*\*cloning\*\*\* of transglutaminase from *Streptovorticillum cinamomeum* CBS 663.68

TI Cloning and expression of natural and synthetic genes for a transglutaminase  
IN Takegi, Hiroshi; Araiuka, Shiro; Matsui, Hiroshi; Washizu, Kinya; Ando, Keiichi; Koikeda, Satoshi  
PA Amano Pharmaceutical Co., Ltd., Japan; Ajinomoto Co., Inc.  
SO Eur. Pat. Appl., 55 pp. CODEN: EPXXDW

DT Patent LA English  
FAN/CNT 1 PATENT NO. KIND DATE APPLICATION NO. DATE \_\_\_\_\_  
PI EP 481504 A1 19920422 EP 91-117813 19911018 EP 481504 B1 19960117 R. DE, FR, GB, JP  
US 1998383 A2 19930810 JP 91-267860 19911016 US 5420025 A 19950530 US 93-136993 19931018  
PRI JP 90-282566 19901019 US 91-777447 19911018  
AB Genes for a transglutaminase useful in food processing and modification of proteins are cloned and expressed. The gene for a transglutaminase of a *Streptovorticillum* was cloned from a BamHI partial digest bank in  $\lambda$ EMBL3 using a probe prep. by polymerase chain reaction amplification of part of the gene using amino acid sequence-derived oligonucleotide primers. Synthetic genes with codon usage optimized for different hosts were prepd. One such gene was expressed in *Escherichia coli* using the ompA-based expression cassette of pIN-III-ompA2. The gene was expressed upon induction with IPTG with most of the transglutaminase activity found in the periplasm. Expression of the gene in yeast and other *Actinomyces* is also demonstrated.

TI Epidermal differentiation and squamous metaplasia: from stem cell to cell death

L6 ANSWER 26 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Tissue transglutaminase cDNA of human erythroleukemia cells

L6 ANSWER 27 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Down-regulated proteins of mesenchymal tumor cells

L6 ANSWER 28 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI \*\*\*Cloning\*\*\* and gene sequence of novel endoglycanases from *Calbio* mixus and *C. gluxus*

L6 ANSWER 29 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Expression of retinoic acid, tritrocholine, and glucocorticoid hormone nuclear receptors is decreased in the liver of rats fed a hypercholesterolemia-inducing diet

L6 ANSWER 30 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Genetic and immunohistochemical detection of mutations inactivating the keratinocyte transglutaminase in patients with lamellar ichthyosis

L6 ANSWER 31 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Method for producing baked goods with improved freshness by using amylase

L6 ANSWER 32 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Isolation of a cDNA encoding a novel member of the transglutaminase gene family from human keratinocytes. Detection and identification of transglutaminase gene products based on reverse transcription-polymerase chain reaction with degenerate primers

L6 ANSWER 33 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Prenatal exclusion of lamellar ichthyosis based on identification of two new mutations in the transglutaminase 1 gene

L6 ANSWER 34 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Defective stratum corneum and early neonatal death in mice lacking the gene for transglutaminase 1 (keratinocyte transglutaminase)

L6 ANSWER 35 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Polyma middle T antigen in HL-60 cells accelerates hematopoietic myeloid and monocytic cell differentiation

L6 ANSWER 36 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI An Egr6-like protein from the filarial parasite *Drillifilaria immitis* has both transglutaminase and protein disulfide isomerase activity

L6 ANSWER 37 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Lessons to learn from the cell death and heat shock \*\*\*genes\*\*\* of *Caenorhabditis elegans*

L6 ANSWER 38 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Effect of AP-1 transcription factors on the regulation of transcription in normal human epidermal keratinocytes

L6 ANSWER 39 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Activation of the human transglutaminase 1 promoter in transgenic mice: terminal differentiation-specific expression of the TGM1-lacZ transgene in keratinized stratified squamous epithelia

L6 ANSWER 40 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI The promoter of the mouse tissue transglutaminase gene directs tissue-specific, retinoid-regulated and apoptosis-linked expression

L6 ANSWER 41 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Transglutaminase-catalyzed inactivation of glyceraldhyde 3-phosphate dehydrogenase and  $\alpha$ -ketoglutarate dehydrogenase complex by polyglutamine domains of pathological length

L6 ANSWER 42 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI The complete genome sequence of the gram-positive bacterium *Bacillus subtilis*

L6 ANSWER 43 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Three novel point mutations in the keratinocyte transglutaminase (TGM) gene in lamellar ichthyosis: significance for mutant transcript level, TGM immunodetection and activity

L6 ANSWER 44 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Reduced expression of tissue transglutaminase in a human endothelial cell line leads to changes in cell spreading, cell adhesion and reduced polymerization of fibronectin

L6 ANSWER 45 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Aging decreases retinoic acid and tritrocholine nuclear expression in rat liver: exogenous retinoid and retinoic acid differentially modulate this decreased expression

L6 ANSWER 46 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Isolation and characterization of a potential gene coding for transglutaminase in *Medicago sativa* L. (alfalfa)

L6 ANSWER 47 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Transglutaminase 1 mutations in autosomal recessive congenital ichthyosis: private and recurrent mutations in an isolated population

L6 ANSWER 48 OF 208 CAPLUS COPYRIGHT 1998 ACS  
TI Biologically Active Heteroantibodies Exhibiting Anticancer Activity and Decreased Toxicity

L6 ANSWER 49 OF 208 CAPLUS COPYRIGHT 1998 ACS

- T1 Tissue transglutaminase-dependent posttranslational modification of the retinoblastoma gene product in promonocytic cells undergoing apoptosis
- L6 ANSWER 50 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Organization and structure of the human tissue transglutaminase gene
- L6 ANSWER 51 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Analysis of the *Bacillus subtilis* genome: \*\*\*cloning\*\*\* and nucleotide sequence of a 62 kb region between 275.degree. (rmb) and 284.degree. (pae)
- L6 ANSWER 52 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Opposite effects of the acute promyelocytic leukemia PML-retinoic acid receptor, -alpha. (RAR.alpha.) and PLZF-RAR.alpha. fusion proteins on retinoic acid signaling
- L6 ANSWER 53 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Age-related decreases in mRNA for brain nuclear receptors and target \*\*\*genes\*\*\* are reversed by retinoic acid treatment
- L6 ANSWER 54 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Antiproliferative effect of curcumin (diferuloylmethane) against human breast tumor cell lines
- L6 ANSWER 55 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Polyglutamine domains are substrates of tissue transglutaminase: does transglutaminase play a role in expanded CAG/PolQ-Q neurodegenerative diseases?
- L6 ANSWER 56 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 High-level expression of the chemically synthesized gene for microbial transglutaminase from *Streptococcus* in *Escherichia coli*
- L6 ANSWER 57 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 A new transglutaminase-like from the ascidian *Ciona intestinalis*
- L6 ANSWER 58 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 T(Gase/C.alpha) protein expression inhibits adenylate cyclase activity in Balb-C 3T3 fibroblasts membranes
- L6 ANSWER 59 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Regulation of the expression of the tissue transglutaminase gene by DNA methylation
- L6 ANSWER 60 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Analysis and comparison of partial sequences of \*\*\*clones\*\*\* from a taste-tud enriched cDNA library
- L6 ANSWER 61 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Growth suppression of squamous cell carcinoma cell lines by PKCs. Possible application to gene therapy
- L6 ANSWER 62 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Phase display and catcher molecules (e.g., suicide inhibitors or transition state analogs) for screening detergent enzyme variants
- L6 ANSWER 63 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Expression of GTP-dependent and GTP-independent tissue-type transglutaminase in cytokine-treated rat brain astrocytes. [Erratum to document cited in CA126:223254]
- L6 ANSWER 64 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 A role for tissue transglutaminase in hepatic injury and fibrogenesis, and its regulation by NF-kappa B
- L6 ANSWER 65 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Tissue transglutaminase in mesenchymal tumor cells
- L6 ANSWER 66 OF 208 CAPLUS COPYRIGHT 1998 ACS
- T1 Expression of GTP-dependent and GTP-independent tissue-type transglutaminase in cytokine-treated rat brain astrocytes
- L6 ANSWER 67 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 alpha 1-adrenergic receptor signaling via Gq is subtype specific and independent of its transglutaminase activity
- L6 ANSWER 68 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Transcription factor regulation of epidermal keratinocyte gene expression
- L6 ANSWER 69 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Efficient production of soluble transglutaminase through co-transformation of *Escherichia coli* with heat shock protein DnaI
- L6 ANSWER 70 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Negative regulation of two hyperproliferative keratinocyte differentiation markers by a retinoic acid receptor-specific retinoid: insight into the mechanism of retinoid action in psoriasis
- L6 ANSWER 71 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 \*\*\*Cloning\*\*\* and sequence analysis of a cDNA encoding salmon (*Oncorhynchus teta*) liver transglutaminase
- L6 ANSWER 72 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Mol. screening and PCR \*\*\*cloning\*\*\* of novel endoglycanases from fungi for use as detergents, textile treatment, and paper pulp processing
- L6 ANSWER 73 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Transglutaminase induction by various cell death and apoptosis pathways
- L6 ANSWER 74 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 *Bacillus*-derived transglutaminase gene sequence, recombinant enzyme production, and use in food industry for crosslinked protein production
- L6 ANSWER 75 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Composition for introducing nucleic acid complexes into higher eukaryotic cells
- L6 ANSWER 76 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Transglutaminases from *Comptosia*, their production with recombinant cells, and their use in foods and cosmetics
- L6 ANSWER 77 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Expression of human transglutaminase C (Tgase II) in yeast *Saccharomyces cerevisiae*: an effect of a domain from carboxyl terminal deletion of the enzyme
- L6 ANSWER 78 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Analysis of up-regulated \*\*\*genes\*\*\* during chondrocyte hypertrophy
- L6 ANSWER 79 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Identification and characterization of up-regulated \*\*\*genes\*\*\* during chondrocyte hypertrophy
- L6 ANSWER 80 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Regulation of gene expression during squamous differentiation by multiple retinoic acid signaling pathways
- L6 ANSWER 81 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Discovery of a new type of proteinase inhibitor family whose members have an anchoring sequence
- L6 ANSWER 82 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Tripeptidyl aminopeptidases from *Aspergillus niger* and *A. oryzae*
- L6 ANSWER 83 OF 208 CAPLUS COPYRIGHT 1998 ACS
- T1 Retinoic acid induction of the tissue transglutaminase promoter is mediated by a novel response element
- L6 ANSWER 84 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 An enzyme and enzyme preparation with endoglycanase activity from *Acartonium* species
- L6 ANSWER 85 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Human prostatic or placental transglutaminase cDNA sequences, recombinant enzyme production, and uses of transglutaminases
- L6 ANSWER 86 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Microbial transglutaminases, their production, gene \*\*\*cloning\*\*\* and sequence, and use for protein crosslinking
- L6 ANSWER 87 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Tissue-specific and androgen-regulated expression of human prostate-specific transglutaminase
- L6 ANSWER 88 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Squamous carcinoma cell lines fail to respond to 1,25-dihydroxyvitamin D despite normal levels of the vitamin D receptor
- L6 ANSWER 89 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Retinoid-induced differentiation of acute promyelocytic leukemia involves PML-RAR.alpha.-mediated increase of type II transglutaminase
- L6 ANSWER 90 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 The proximal promoter of the human transglutaminase 3 gene, stratified squamous epithelial-specific expression in cultured cells is mediated by binding of Sp1 and ets transcription factors to a proximal promoter element
- L6 ANSWER 91 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 \*\*\*Cloning\*\*\* and regulation of corfilin, beta., a new member of the corfilin/pr family. Suppression by retinoic acid receptor-selective retinoids
- L6 ANSWER 92 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Subtyping of coagulation factor XIIIa
- L6 ANSWER 93 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Biochemical, structural, and transglutaminase substrate properties of human loricin, the major epidermal cornified cell envelope protein
- L6 ANSWER 94 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Transglutaminase originating in Japanese oyster
- L6 ANSWER 95 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Involvement of retinoic acid nuclear receptors in transcriptional regulation of tissue transglutaminase, the gene involved in apoptosis
- L6 ANSWER 96 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Tissue-type transglutaminase from red sea bream (*Pagrus major*). Sequence analysis of the cDNA and functional expression in *Escherichia coli*
- L6 ANSWER 97 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 \*\*\*Cloning\*\*\*, characterization, and tissue distribution of porcine SPAL, a protein with a transglutaminase substrate domain and the WAP motif
- L6 ANSWER 98 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 The importance of the GTP-binding protein tissue transglutaminase in the regulation of cell cycle progression
- L6 ANSWER 99 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Autosomal recessive lamellar ichthyosis: identification of a new mutation in transglutaminase 1 and evidence for genetic heterogeneity
- L6 ANSWER 100 OF 208 CAPLUS COPYRIGHT 1998 ACS
- T1 Localization of the human prostatic transglutaminase (Type IV) gene (TGM4) to chromosome 3p21.33-p22 by fluorescence in situ hybridization
- L6 ANSWER 101 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Use of heparanase to identify and isolate anti-heparanase compound
- L6 ANSWER 102 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Molecular \*\*\*cloning\*\*\* and expression of cDNA for fish transglutaminase
- L6 ANSWER 103 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Isolation and characterization of the human tissue transglutaminase gene promoter
- L6 ANSWER 104 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 \*\*\*Genes\*\*\* up-regulated in hypertrophied ventricle
- L6 ANSWER 105 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Proliferation-dependent vs. independent programmed cell death of prostatic cancer cells involves distinct gene regulation
- L6 ANSWER 106 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Fermentative production of active enzyme with proenzyme-producing cells in proenzyme-cleaving protease-containing medium
- L6 ANSWER 107 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Alterations in murine keratinocyte differentiation induced by activated rasHa \*\*\*genes\*\*\* are mediated by protein kinase C-alpha.
- L6 ANSWER 108 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Restoration of differentiation and suppression of tumorigenicity in somatic cell hybrids of human squamous carcinoma cells and keratinocytes
- L6 ANSWER 109 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Assignment of the human transglutaminase 2 (TGM2) and transglutaminase 3 (TGM3) \*\*\*genes\*\*\* to chromosome 20q11.2
- L6 ANSWER 110 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Structure and organization of the human transglutaminase 3 gene: evolutionary relationship to the transglutaminase family
- L6 ANSWER 111 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Molecular \*\*\*cloning\*\*\* of mouse erythrocyte protein 4.2, a membrane protein with strong homology with the transglutaminase superegene family
- L6 ANSWER 112 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 The structure of microbial transglutaminase from *Streptococcus* and its gene expression in *Escherichia coli*
- L6 ANSWER 113 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 The human tissue transglutaminase gene maps on chromosome 20q12 by in situ fluorescence hybridization
- L6 ANSWER 114 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Molecular \*\*\*cloning\*\*\* and characterization of a novel transglutaminase cDNA from a human prostate cDNA library
- L6 ANSWER 115 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 \*\*\*cloning\*\*\* reveals domains that regulate its specific activity and substrate specificity
- L6 ANSWER 116 OF 208 CAPLUS COPYRIGHT 1998 ACS  
T1 Cosmetic compositions containing corneocyte proteins
- L6 ANSWER 117 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Transfection of tissue transglutaminase into a highly malignant hamster fibrosarcoma leads to a reduced incidence of primary tumor growth	T1 Transformation of NIH3T3 cells with ras oncogenes abrogates the retinoic acid induction of tissue transglutaminase	L6 ANSWER 126 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Chemical synthesis of the gene for microbial transglutaminase from Streptovorticillum and its expression in Escherichia coli	L6 ANSWER 126 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Wild-type p53 tumor suppressor gene restores differentiation of human squamous carcinoma cells but not the response to transforming growth factor beta.	L6 ANSWER 123 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Transglutaminases: protein crosslinking enzymes in tissues and body fluids	L6 ANSWER 121 OF 208 CAPLUS COPYRIGHT 1998 ACS	L6 ANSWER 120 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Preparation of bacterial transglutaminase with Escherichia coli	L6 ANSWER 118 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Transferrin expression during involution of the rat ventral prostate after castration	L6 ANSWER 119 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Differential responsiveness of human bronchial epithelial cells, lung carcinoma cells, and bronchial fibroblasts to interferon-gamma, in vitro	L6 ANSWER 122 OF 208 CAPLUS COPYRIGHT 1998 ACS	L6 ANSWER 125 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** of DNA encoding mammalian trichoglycin and transglutaminases-3 and use of these proteins for formation of gels for use in food, cosmetics, and medicine	L6 ANSWER 124 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** and characterization of the full length cDNA and promoter of mouse tissue transglutaminase	L6 ANSWER 125 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** of the gene for microbial transglutaminase from Streptovorticillum and its expression in Streptomyces lividans	L6 ANSWER 127 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Molecular ***cloning*** of the gene for microbial transglutaminase	L6 ANSWER 128 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 HIV-1 gp120-dependent induction of apoptosis in antigen-specific human T cell ***clones*** is characterized by 'tissue' transglutaminase expression and prevented by cyclosporin A	L6 ANSWER 129 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Induction of gene expression during involution of the lactating mammary gland of the rat	L6 ANSWER 130 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Control of growth regulatory and differentiation-specific ***genes*** in human epidermal keratinocytes by interferon gamma. Antagonism by retinoic acid and transforming growth factor beta.1	L6 ANSWER 131 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Regulation of proliferation-specific and differentiation-specific ***genes*** during senescence of human epidermal keratinocyte and mammary epithelial cells	L6 ANSWER 132 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Differential gene regulation during programmed death (apoptosis) versus proliferation of prostatic glandular cells induced by androgen manipulation	L6 ANSWER 133 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Transformation of NIH3T3 cells with ras oncogenes abrogates the retinoic acid induction of tissue transglutaminase	L6 ANSWER 134 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Site-directed mutagenesis of human tissue transglutaminase. Cys-277 is essential for transglutaminase activity but not for G1Tase activity	L6 ANSWER 135 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** of the cDNA encoding transglutaminase of fish	L6 ANSWER 136 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** of human prostatic transglutaminases	L6 ANSWER 137 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 The deduced sequence of the novel protransglutaminase E (TGase3) of human and mouse	L6 ANSWER 138 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Human epidermal type I transglutaminase gene promoter and its use in tissue-specific expression of ***genes***	L6 ANSWER 139 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 SKALP/Plcfr: An elastase inhibitor from cultured human keratinocytes. Purification, cDNA sequence, and evidence for transglutaminase cross-linking	L6 ANSWER 140 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Corfmin, a cross-linked envelope precursor in keratinocytes that is down-regulated by retinoids	L6 ANSWER 141 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Differences in the post-translational modification of proteins by polyamines between weakly and highly metastatic B16 melanoma cells	L6 ANSWER 142 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** and expression of chicken erythrocyte transglutaminase	L6 ANSWER 143 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Identification of Gln26 in nitrogen as the amine acceptor in transglutaminase-catalyzed cross-linking of laminin-nitrogen complexes	L6 ANSWER 144 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Urmus hemocyte transglutaminase. cDNA ***cloning***, amino acid sequence, and tissue localization	L6 ANSWER 145 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Identification of promoter region of guinea pig liver transglutaminase gene	L6 ANSWER 146 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** of bovine transglutaminase cDNA	L6 ANSWER 147 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 A retinoic acid-inducible mRNA from human erythroleukemia cells encodes a novel tissue transglutaminase homolog	L6 ANSWER 148 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** of human epidermal transglutaminase for recombinant manufacture of the enzyme	L6 ANSWER 149 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Multiple cell cycle access to the apoptotic death program in human neuroblastoma cells	L6 ANSWER 150 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Structure of the gene for human epidermal transglutaminase	L6 ANSWER 151 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 The cornified cell envelope: Loricin and transglutaminases	L6 ANSWER 152 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Molecular ***cloning*** of rat prostate transglutaminase complementary DNA. The major androgen-regulated protein DP1 of rat dorsal prostate and coagulating gland	L6 ANSWER 153 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Retinoids and state of differentiation modulate CRABP II gene expression in a skin equivalent	L6 ANSWER 154 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Genomic structure of keratinocyte transglutaminase. Recruitment of new exon for modified function	L6 ANSWER 155 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** and sequence analysis of cDNA ***clones*** for bovine aortic endothelial cell transglutaminase	L6 ANSWER 156 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Phenotype-specific 'tissue' transglutaminase regulation in human neuroblastoma cells in response to retinoic acid: correlation with cell death by apoptosis	L6 ANSWER 157 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** and expression of ***genes*** for human and mouse tissue transglutaminase	L6 ANSWER 158 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** and expression of natural and synthetic ***genes*** for a transglutaminase	L6 ANSWER 159 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Coupling of adenovirus to transferin-polyvinylDNA complexes greatly enhances receptor-mediated gene delivery and expression of transfected ***genes***	L6 ANSWER 160 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Type I/keratinocyte transglutaminase: expression in human skin and psoriasis	L6 ANSWER 161 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Polyphosphates containing the fibrin-binding domain of fibronectin, their recombinant production, and their use in imaging and therapy	L6 ANSWER 162 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Structure and organization of the human transglutaminase 1 gene	L6 ANSWER 163 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Organization of the gene for human erythrocyte membrane protein 4.2: structural similarities with the gene for a subunit of factor XIII	L6 ANSWER 164 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Expression of keratinocyte transglutaminase mRNA revealed by in situ hybridization	L6 ANSWER 165 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Recombinant coagulation factor XIII/XIIIa, mutant analogs thereof, and use in imaging of blood clots	L6 ANSWER 166 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Induction of peroxisomal beta-oxidation ***genes*** by retinoic acid in cultured rat hepatocytes	L6 ANSWER 167 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Blood-coagulation factor XIII manufacture with recombinant yeast	L6 ANSWER 168 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Expression of the transglutaminase gene in Escherichia coli	L6 ANSWER 169 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Molecular ***cloning*** of human epidermal transglutaminase cDNA from keratinocytes in culture	L6 ANSWER 170 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Process for the expression of foreign ***genes*** in yeasts	L6 ANSWER 171 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 The expression of 'tissue' transglutaminase in two human cancer cell lines is related with the programmed cell death (apoptosis)	L6 ANSWER 172 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Epidermal type I transglutaminase (TGM1) is assigned to human chromosome 14	L6 ANSWER 173 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Primary structure of keratinocyte transglutaminase	L6 ANSWER 174 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Isolation and characterization of cDNA ***clones*** to mouse macrophage and human endothelial cell tissue transglutaminases	L6 ANSWER 175 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 The complete amino acid sequence of the human transglutaminase K enzyme deduced from the nucleic acid sequences of cDNA ***clones***	L6 ANSWER 176 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Isolation of cDNA for human epidermal type I transglutaminase	L6 ANSWER 177 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** of gene for human fibronectin analogs: its recombinant manufacture, and pharmaceuticals contg. same	L6 ANSWER 178 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Expression vectors for synthesis of heterologous proteins in Schizosaccharomyces pombe	L6 ANSWER 179 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Differentiation capacity of human non-small-cell lung cancer cell lines after exposure to photod ester	L6 ANSWER 180 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 ***Cloning*** of mammalian type I transglutaminase cDNA	L6 ANSWER 181 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Molecular nature of in vivo mutations in human cells at the autosomal HLA-A locus	L6 ANSWER 182 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Complete amino acid sequence and homologies of human erythrocyte membrane protein band 4.2	L6 ANSWER 183 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Expression of functional coagulation factor XIII in Escherichia coli	L6 ANSWER 184 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Molecular ***cloning*** of human protein 4.2: a major component of the erythrocyte membrane	L6 ANSWER 185 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Recombinant manufacture of transglutaminase of Cavidae liver (MTGase) with Escherichia	L6 ANSWER 186 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Histological and biochemical characterization of the murine cataract mutant Nop	L6 ANSWER 187 OF 208 CAPLUS COPYRIGHT 1998 ACS	T1 Deletion and linkage mapping of eight markers from the proximal short arm of chromosome 6	L6 ANSWER 188 OF 208 CAPLUS COPYRIGHT 1998 ACS	
--	---	--	---	--	--	--	---	--	--	--	--	--	--	---	--	--	--	--	---	--	---	--	---	--	---	--	---	--	--	--	---	--	--	--	---	--	---	--	--	--	---	--	--	--	---	--	---	--	--	--	---	--	---	--	--	--	--	--	--	--	--	--	--	--	--	--	---	--	---	--	---	--	--	--	--	--	--	--	---	--	--	--	--	--	---	--	---	--	---	--	---	--	--	--	--	--	---	--	---	--	---	--	---	--	--	--	---	--	--	--	---	--	--	--	---	--	--	--	---	--	--	--	--	--	---	--	---	--	--	--	--	--	---	--	---	--	--	--	---	--	--	--	--	--	--

T1 Production of recombinant eukaryotic cells efficiently expressing recombinant \*\*\*genes\*\*\*

L6 ANSWER 186 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Differential reaction of secretory and non-secretory proteins in hormone-treated burning tumor

L6 ANSWER 190 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Regulation of type I (epidermal) transglutaminase mRNA levels during squamous differentiation: down regulation by retinoids

L6 ANSWER 191 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Fibrin-binding peptides and method for detection of fibrin deposits

L6 ANSWER 192 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 New expression vectors for the fission yeast Schizosaccharomyces pombe

L6 ANSWER 193 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Use of transglutaminase

L6 ANSWER 126 OF 208 CAPLUS COPYRIGHT 1998 ACS AN 1994:209639 CAPLUS DN 120:209639

T1 Chemical synthesis of the gene for microbial transglutaminase from Streptovorticillum and its expression in Escherichia coli

AU Takehana, Shinjo, Washizu, Kinya, Ando, Keiichi, Koikeda, Satoshi, Takeuchi, Kazuyuki, Matsui, Hiroshi, Motoki, Masao, Takagi, Hiroshi

CS Food Res. Dev. Lab., Ajinomoto Co., Inc., Kawasaki, 210, Japan

SO Biosci. Biotechnol., Biochem. (1994), 58(1), 88-92 CODEN: BBBI EJ, ISSN: 0916-8451

DI Journal LA English

AB The gene coding for microbial transglutaminase (TGase) from Streptovorticillum, which consists of 331 amino acids, was chem. synthesized. The codons have been substituted for those mainly favored in yeast. The authors' strategy involved the construction of the TGase gene in five sections (34 oligomers) that contained unique restriction enzyme sites at both ends, which could readily be ligated to form the full-length product. The chem. synthesized gene was inserted downstream from the ompA signal peptide of the E. coli expression vector, pJN-III-ompA, which carries lpp and lac promoters. The resultant plasmid directed the expression of TGase, with the activity being secreted mainly into the periplasmic space of E. coli. The induced gene product was identical with native TGase in size and in immunol. properties, though the enzyme activity was low.

L6 ANSWER 127 OF 208 CAPLUS COPYRIGHT 1998 ACS AN 1994:209638 CAPLUS DN 120:209638

T1 Molecular \*\*\*cloning\*\*\* of the gene for microbial transglutaminase from Streptovorticillum and its expression in Streptomyces lividans

AU Washizu, Kinya, Ando, Keiichi, Koikeda, Satoshi, Hirose, Susumu, Matsura, Akira, Takagi, Hiroshi, Motoki, Masao, Takeuchi, Kazuyuki

CS Tsukuba Res. Lab., Ariano Pharm. Co., Ltd., Tsukuba, 305, Japan

SO Biosci. Biotechnol., Biochem. (1994), 58(1), 82-7 CODEN: BBBI EJ, ISSN: 0916-8451

DI Journal LA English

AB The microbial transglutaminase (TGase)-producing strain S-8-112 was identified as a variant of Streptovorticillum mobaraense. The authors amplified a partial gene fragment by polymerase chain reaction (PCR) using oligonucleotides synthesized from the amino acid sequence of TGase, and \*\*\*cloned\*\*\* the gene for TGase using the PCR amplified fragment as a probe. The gene encoded a precursor of TGase consisting of 406 amino acid residues, which comprised the prepro region of 75 amino acid residues and the mature region of 331 amino acid residues. The authors expressed the TGase gene in Streptomyces lividans under a tyrosinase promoter, and found an active and mature recombinant enzyme, indicating the processing of the gene product.

L6 ANSWER 185 OF 208 CAPLUS COPYRIGHT 1998 ACS AN 1990:438931 CAPLUS DN 113:38931

T1 Recombinant manufacture of transglutaminase of Cavitation liver (MTGase) with Escherichia

IN Ikura, Koji, Sasaki, Ryuzo, Chiba, Hideo

PA Ajinomoto Co., Inc., Japan

SO Jpn. Kokai Tokkyo Koho, 8 pp. CODEN: JKKXAF

DI Patent LA Japanese

FAN/CNT 1 PATENT NO. KIND DATE APPLICATION NO. DATE \_\_\_\_\_

PI JP 01300889 A2 19891205 JP 88-132000 19880530

AB A method for manu. MTGase by cultivating recombinant E. coli is described. cDNA for MTGase was \*\*\*cloned\*\*\* from a guinea pig liver cDNA library and subsequently used to construct an expression plasmid pK1G1. The E. coli transformants were cultured and induced to produce MTGase died by Western blotting. Purif. of the recombinant MTGase with monoclonal antibody to MTGase by affinity chromatog. was given. The purified MTGase had a sp. activity of 1690 unit/mg times. 104.

L6 ANSWER 199 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 \*\*\*Cloning\*\*\* of cDNA coding for guinea pig liver transglutaminase

L6 ANSWER 200 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Characterization of cDNA coding for human factor XIIIa

L6 ANSWER 201 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Regulation of proliferation and differentiation of respiratory tract epithelial cells by TGF-beta

L6 ANSWER 202 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Amino acid sequence of the a subunit of human factor XIII

L6 ANSWER 203 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Transglutaminase activity and putrescine-binding capacity in \*\*\*cloned\*\*\* cell lines with different metastatic potential

L6 ANSWER 204 OF 208 CAPLUS COPYRIGHT 1998 ACS

L6 ANSWER 194 OF 208 CAPLUS COPYRIGHT 1998 ACS AN 1988:623906 CAPLUS DN 109:223906

T1 Improved "ATG vector" series for bacterial synthesis of proteins and protein fragments

AU Broeker, Michael

CS Res. Lab., Behringwerke A.-G., Marburg D-3550, Fed. Rep. Ger.

SO Biotechniques (1988), 6(8), 734 CODEN: BTNQDQ, ISSN: 0735-6205

DI Journal LA English

AB The ATG vector developed by E. Anann and J. Brosius (1985) facilitates regulated high level expression of \*\*\*genes\*\*\* in Escherichia coli. This vector was improved by insertion of a multiple \*\*\*cloning\*\*\* site from pUC12 downstream of the translational start codon, and offers in-frame ligation in at 3 potential reading frames of coding DNA. This improved series of POB vectors was shown to direct the efficient expression of coding sequences of lacZ and the heat labile enterotoxin subunit B of E. coli and of the human blood coagulation protein XIIIa.

L8 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1998 ACS AN 1994:209639 CAPLUS DN 120:209639

T1 Chemical synthesis of the gene for microbial transglutaminase from Streptovorticillum and its expression in Escherichia coli

AU Takehana, Shinjo, Washizu, Kinya, Ando, Keiichi, Koikeda, Satoshi, Takeuchi, Kazuyuki, Matsui, Hiroshi, Motoki, Masao, Takagi, Hiroshi

CS Food Res. Dev. Lab., Ajinomoto Co., Inc., Kawasaki, 210, Japan

SO Biosci. Biotechnol., Biochem. (1994), 58(1), 88-92 CODEN: BBBI EJ, ISSN: 0916-8451

DI Journal LA English

AB The gene coding for microbial transglutaminase (TGase) from Streptovorticillum, which consists of 331 amino acids, was chem. synthesized. The \*\*\*codons\*\*\* have been \*\*\*substituted\*\*\* for those mainly favored in yeast. The authors' strategy involved the construction of the TGase gene in five sections (34 oligomers) that contained unique restriction enzyme sites at both ends, which could readily be ligated to form the full-length product. The chem. synthesized gene was inserted downstream from the ompA signal peptide of the E. coli expression vector, pJN-III-ompA, which carries lpp and lac promoters. The resultant plasmid directed the expression of TGase, with the activity being secreted mainly into the periplasmic space of E. coli. The induced gene product was identical with native TGase in size and in immunol. properties, though the enzyme activity was low.

T1 Variable transglutaminase activity in human diploid fibroblasts during in vitro senescence

L6 ANSWER 205 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Production of monoclonal antibodies to guinea pig liver transglutaminase

L6 ANSWER 206 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Monoclonal antibody to the region of fibronectin involved in crosslinking to human fibrin

L6 ANSWER 207 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Primary amines inhibit the triggering of B lymphocytes to antibody synthesis

L6 ANSWER 208 OF 208 CAPLUS COPYRIGHT 1998 ACS

T1 Lysosomotropic agents modulate serum stimulation of dome formation in MDCK epithelial cell cultures

\*\*\*\*\*

\* \* \* \* \*

\* **Welcome to MESSENGER (APS Text) at USPTO** \*

\* **The USPTO production files are current through:** \*

\* **DECEMBER 15, 1998 for U.S. Patent Text Data.** \*

\* **DECEMBER 15, 1998 for U.S. Current Classification Data.** \*

\* **DECEMBER 15, 1998 for U.S. Patent Image Data.**

FILE 'USPAT' ENTERED AT 13:06:08 ON 21 DEC 1998

(FILE 'USPAT' ENTERED AT 13:06:08 ON 21 DEC 1998)

L1 260 S TRANSGLUTAMINASE?

L2 1146 S CODON?(10N)(OPTIMIZ? OR SUBSTIT?)

L3 8 S L1 AND L2

L4 1 S L1(P)L2

L5 97625 S DELETF



## L6 5 L5(P)L1

- L3, 1, 5,817,768, Oct. 6, 1998, Monospecific antibodies against a subunit of fibrinogen; Gerd Grieninger, et al., 530388, 1; 435/7, 13, 530388,25, 389,3, 391,1, 391,3 [IMAGE AVAILABLE]
- 2, 5,782,742, Aug. 11, 1998, Fibrin-binding peptide fragments of fibrinectin; Leslie I. Gold, et al., 514/2, 424/9, 1; 435/69,6, 514/8; 530350, 402 [IMAGE AVAILABLE]
- 3, 5,616,500, Apr. 1, 1997, Trichoyalin and "transglutaminase"-3 and methods of using same; Peter M. Steinert, et al., 435/320, 1, 69, 1, 193, 325, 348; 536/22, 1, 23, 1, 23, 2, 23, 5, 24, 3, 24, 33 [IMAGE AVAILABLE]
- 4, 5,527,692, Jun. 18, 1996, Methods for producing thrombin; Richard D. Holly, et al., 424/94,64; 435/69, 1, 69, 6, 214, 320, 1; 514/2, 12; 530350, 380, 361, 382; 536/22, 1, 23, 1, 23, 2, 23, 5 [IMAGE AVAILABLE]
- 5, 5,514,579, May 7, 1996, Human "transglutaminases"; Patrick J. O'Hara, et al., 435/352, 69, 2, 193, 254, 3, 320, 1; 536/23, 2, 24, 31 [IMAGE AVAILABLE]
- 6, 5,502,034, Mar. 26, 1996, Methods for producing thrombin; Richard D. Holly, et al., 514/12; 424/94,64; 435/69, 1, 69, 6, 214, 320, 1; 514/2; 530350, 380, 381, 382; 536/22, 1, 23, 1, 23, 2, 23, 5 [IMAGE AVAILABLE]
- 7, 5,486,599, Jan. 23, 1996, Construction and use of synthetic constructs encoding syndecan; Scott Saunders, et al., 530395; 435/69, 1, 69, 7, 252, 3, 320, 1; 536/23, 4, 23, 5 [IMAGE AVAILABLE]
- 8, 5,476,777, Dec. 19, 1995, Methods for producing thrombin; Richard D. Holly, et al., 435/214; 424/94,64; 435/69, 1, 69, 6, 254, 11, 254, 2, 254, 21, 320, 1, 352; 530350, 380, 381, 382; 536/22, 1, 23, 1, 23, 2, 23, 5 [IMAGE AVAILABLE]
- L4, 1, 5,514,579, May 7, 1996, Human transglutaminases; Patrick J. O'Hara, et al., 435/352, 69, 2, 193, 254, 3, 320, 1; 536/23, 2, 24, 31 [IMAGE AVAILABLE]

US PAT NO: 5,514,579 [IMAGE AVAILABLE] L6: 1 of 1  
BSJUM(2) As . . . the art, the DNA molecules of the present invention encompass allelic variants and genetically engineered or synthetic variants of the "transglutaminases" that encode conservative amino acid substitutions and/or minor additions, or deletions of amino acids. Such variants also encompass DNA molecules containing degeneracies in the DNA code wherein host-preferred "codons" are "substituted" for the analogous "codons" in the human sequence. In addition, substantially similar DNA molecules of the present invention encompass those DNA molecules that are . . .

- L6, 1, 5,712,252, Jan. 27, 1998, Method of augmenting soft tissue in mammals; Dean Preston Smith, 514/21, 424/423, 424 [IMAGE AVAILABLE]
- 2, 5,607,849, Mar. 4, 1997, Gene encoding transglutaminase derived from fish; Hisashi Yasueda, et al., 435/193, 69, 1, 183, 252,31, 252,33, 254, 11, 254, 21, 320, 1; 536/23, 2 [IMAGE AVAILABLE]
- 3, 5,514,579, May 7, 1996, Human transglutaminases; Patrick J. O'Hara, et al., 435/352, 69, 2, 193, 254, 3, 320, 1; 536/23, 2, 24, 31 [IMAGE AVAILABLE]
- 4, 5,514,573, May 7, 1996, Gene encoding transglutaminase derived from fish; Hisashi Yasueda, et al., 435/193, 69, 1, 183, 320, 1; 536/23, 2 [IMAGE AVAILABLE]
- 5, 5,428,014, Jun. 27, 1995, Transglutaminase cross-linkable polypeptides and methods relating thereto; Virender Labroo, et al., 514/12, 13, 14, 15, 16; 530324, 326, 327, 328, 329, 345, 350 [IMAGE AVAILABLE]

US PAT NO: 5,712,252 [IMAGE AVAILABLE] L6: 1 of 5  
DETD(1) Lorcin . . . Sci U.S.A., 89,910-14 (1992); and Kogge, et al., "The Two Size Alleles of Human Keratin 1 are Due to a "Deletion" in the Glycine-Rich Carboxyl-Terminal V2 Subdomain", J. Invest. Dermatol., 99:697-702 (1992), which are hereby incorporated by reference). These sequences may . . . and promote nonspecific interaction between K1fs and CE. Lorcin incorporation into the cell envelope or cornified envelope (CE) by epidermal "transglutaminases" has been documented in vivo, by identification of lorcin polypeptides directly cross-linked to the CE by isopeptide bonds (FIG. 2).

US PAT NO: 5,607,849 [IMAGE AVAILABLE] L6: 2 of 5  
DETD(18) The present DNA fragment includes mutants having substitution, "deletion" or insertion of base sequences on the basis of the difference in the individualities of fishes and of the difference. . . for example, which may be a pseudogene. However, such still contain an essentially equivalent DNA fragment capable of expressing the "transglutaminase" activity. The presence of them is described in the following examples.

DETD(54) From the above, it was clarified that the "transglutaminase" of SEQ ID NO:7 is an Alaska pollock "transglutaminase" as expressed beyond the kind of the organ, and that the "transglutaminase" of SEQ ID NO:28, though not obtained as a cDNA of a complete length, was different from the liver-derived "transglutaminase" only in the point of a several-base substitution, a base "deletion" of 12 bp and a base insertion of 3 bp in the structural gene. Thus, both genes were clarified to. . .

US PAT NO: 5,514,579 [IMAGE AVAILABLE] L6: 3 of 5  
BSJUM(12) As . . . the art, the DNA molecules of the present invention encompass allelic variants and genetically engineered or synthetic variants of the "transglutaminases" that encode conservative amino acid substitutions and/or minor additions, or "deletions" of amino acids. Such variants also encompass DNA molecules containing degeneracies in the DNA code wherein host-preferred codons are substituted. . .

BSJUM(13) Recombinant DNA expression systems provide convenient means for obtaining large quantities of the human "transglutaminases" in relatively pure form. By human prostatic or placental "transglutaminase" polypeptides and fragments is meant to include sequences of amino acids from 9 to 20 amino acids up to entire. . . more preferably at least about 95% or more homology to the amino acid sequences of the human prostatic or placental "transglutaminases" of the invention. As will be appreciated by those skilled in the art, the invention also includes those polypeptides having. . . genetic polymorphism) or may be produced by human intervention (e.g., by mutagenesis of cloned DNA sequences), such as induced point, "deletion" and insertion mutations.

DETD(19) The prostate "transglutaminase" cDNA insert present in plasmid pDT 4/-15 was subcloned into the mammalian expression vector Zerm229R. Plasmid Zerm229 is a pUC18-based expression. . . and an expression unit containing the SV40 early promoter, mouse dihydropyridine reductase gene, and SV40 terminator. Zerm229 was modified to "delete" the two Eco RI sites by partial digestion with Eco RI, blunting with DNA polymerase I (Klenow fragment) and dNTPs. . .

US PAT NO: 5,514,573 [IMAGE AVAILABLE] L6: 4 of 5  
DETD(18) The present DNA fragment includes mutants having substitution, "deletion" or insertion of base sequences on the basis of the difference in the individualities of fishes and of the difference. . . for example, which may be a pseudogene. However, such still contain an essentially equivalent DNA fragment capable of expressing the "transglutaminase" activity. The presence of them is described in the following examples.

DETD(54) From the above, it was clarified that the "transglutaminase" of SEQ ID NO:7 is an Alaska pollock "transglutaminase" as expressed beyond the kind of the organ, and that the "transglutaminase" of SEQ ID NO:28, though not obtained as a cDNA of a complete length, was different from the liver-derived "transglutaminase" only in the point of a several-base substitution, a base "deletion" of 12 bp and a base insertion of 3 bp in the structural gene. Thus, both genes were clarified to. . .

US PAT NO: 5,428,014 [IMAGE AVAILABLE] L6: 5 of 5  
BSJUM(2) The . . . polypeptides can also be synthesized by conventional solution phase methodology. The polypeptides are then screened for the ability to form "transglutaminase"-induced cross-links using methods described in more detail below. As will be evident to one skilled in the art, polypeptides can be prepared in which the sequence and content of the spacer and flanking sequences can be altered by "deletion", addition or replacement to improve the cross-linking rate and/or to reduce the deamination of the intermediate. For example, the spacer. . . amino acid 7, to Gly, amino acid 12, in the polypeptide sequence Thr-Ile-Gly-Glu-Gly-Gln-His-His-Ile-Gly-Gly-Ala-Lys-Gln-Ala-Lys-Val (SEQ ID NO:1) can be reduced by "deletion" of one or all of the amino acid residues. In a like manner, the amino-terminal sequences flanking Gln, amino acid residue 6, and the carboxy-terminal sequences flanking Lys, amino acid residue 14, can be "deleted" to shorten the polypeptide. It is preferable to place a glycine or proline residue at the carboxyl terminus of the. . .